

# Geometrisation of electromagnetic field and topological interpretation of quantum formalism

O.A.Olkhov

*N.N.Semenov Institute of Chemical Physics, Russian Academy of Sciences*

*Moscow, Russia, E-mail: olkhov@center.chph.ras.ru*

All numerous attempts of the electromagnetic field geometrisation were based on concept "particle-field" (pointlike sources and extended field; see for example [1-3]). But we had recently shown that the free Dirac field can be considered as a curved nonorientable closed connected time-space 4-manifold [4]. Its fundamental group consists of four global gliding symmetries and the Minkowski space appears as the manifold universal covering space. Taking this into account we now suggest a new approach to the electromagnetic field geometrisation and this approach inevitably means a new topological interpretation of the quantum mechanics mathematical formalism.

Namely, we suppose that interacting Dirac and Maxwell fields can be considered as a single closed connected curved 4-manifold. Its fundamental group consists of four local gliding symmetries and its covering space has a more complex curved structure. Electric and magnetic fields appear within such approach as components of the curvature tensor of the manifold covering space and the Dirac spinors appear as basic functions of the manifold fundamental group representation.

Above concept differs from the one of general theory of relativity by two main points: we geometrize not only the field but we also geometrize the field sources and we represent the field and the sources not as a curved riemannian 4-space but as a 4-manifold.

We start with the known equations for interacting classical electromagnetic and electron-positron fields [5]

$$i\gamma_1\left(\frac{\partial}{\partial x_1} + ieA_1\right)\psi - \sum_{\alpha=2}^4 i\gamma_\alpha\left(\frac{\partial}{\partial x_\alpha} + ieA_\alpha\right)\psi = m\psi, \quad (1)$$

$$F_{kl} = \frac{\partial A_k}{\partial x_l} - \frac{\partial A_l}{\partial x_k}, \quad (2)$$

$$\sum_{i=1}^4 \frac{\partial F_{ik}}{\partial x_i} = j_k, \quad j_k = e\psi * \gamma_1\gamma_k\psi. \quad (3)$$

Here  $\hbar = c = 1$ ,  $x_1 = t$ ,  $x_2 = x$ ,  $x_3 = y$ ,  $x_4 = z$ ,  $F_{kl}$ -tensor of electric and magnetic fields,  $A_k$  - 4-potential,  $\gamma_k$ -Dirac matrices,  $\psi$ - Dirac spinor,  $m$  and  $e$ -mass and charge of an electron.

It is shown that in (1) the expression  $\nabla_k = \partial/\partial x_k + ieA_k$  can be considered as the translation group generator into a conformal pseudo-euclidean 4-space and  $ieA_k$  appear within such approach as  $\Gamma_k$  —the contraction of a riemannian connection  $\Gamma_{kp}^q$  of this space

$$\Gamma_k = \sum_p \Gamma_{kp}^p.$$

Multiplied by the reflection operators  $\gamma_k$  [1] the  $\nabla_k$  gives the representation of a local gliding symmetry group in this space with the Dirac spinors as basic vectors of the representation. All this gives the opportunity to interpret (1) as the metric relation for a closed connected nonorientable topological 4-manifold with the local gliding symmetry group as its fundamental group and with a conformal pseudoeuclidean space as its universal covering space (two-dimensional analog of such manifold is the combination of a torus and the Klein bottle [6]).

Now we use a relation between a riemannian connection and a riemannian curvature tensor  $R_{lk,i}^q$  [7]

$$R_{lk,i}^q = \frac{\partial \Gamma_{li}^q}{\partial x_k} - \frac{\partial \Gamma_{ki}^q}{\partial x_l} + \Gamma_{kp}^q \Gamma_{li}^p - \Gamma_{lp}^q \Gamma_{ki}^p.$$

After contraction over  $q$  and  $l$  we obtain

$$R_{ik}^o = \sum_q R_{ik,q}^q = \frac{\partial \Gamma_i}{\partial x_k} - \frac{\partial \Gamma_k}{\partial x_i}. \quad (4)$$

By comparison (2) and (4) we can write down the equations (1-3) using only geometrical notations

$$i\gamma_1 \nabla_1 \psi - \sum_{\alpha=2}^4 i\gamma_\alpha \nabla_\alpha \psi = m\psi, \quad (5)$$

$$\sum_{i=1}^4 \frac{\partial R_{ik}^o}{\partial x_i} = j_k, \quad (6)$$

where  $R_{ik}^o$  has the form (4).

Finally, we have the following geometrical interpretation of electromagnetic field.

1. Electromagnetic field and its sources (electron-positron field) can be considered as a single 4-manifold which fundamental group is generated by four local gliding symmetries.
2. Universal covering space of this manifold is the conformal pseudoeuclidean space.
3. The 4-potentials  $A_k$  is defined by the connections of this space  $\Gamma_k$  ( $ieA_k = \Gamma_k$ ).
4. The electric and magnetic field components are defined by the components of the covering space curvature tensor  $R_{ik}$  ( $ieF_{ik} = R_{ik}$ ).
5. The Dirac spinors appear as basic functions for the manifold fundamental group representation.

One comment in conclusion. We see that replacing a "wave-particle" by a space-time manifold does not mean "more determinism" for the quantum object description. And we also see that the above topological approach does not introduce any hidden variables and does not therefore contradict the Bell and von Neumann theorems [8,9].

1. Weyl H. Gravitation und Electrizitat. Berlin, Sitzber.Preuss.Akad.Wiss, 1918
2. Konopleva N.P., Sokolic H.A., Nucl.Phys. 1965. V.72. P.667
3. Daniel M., Viallet C.M., Rev.Mod,Phys. 1980. V.52. P.175
4. Olkhov O.A., e-print quant-ph/0101137, in Proceedings of the 7th International Symposium on Particles, Strings and Cosmology, Lake Tahoe, California, 10-16 December 1999. Singapore-New Jersey- Hong Kong.: World Scientific, 2000, P.160.

5. Schweber S., An introduction to relativistic quantum field theory, Row, Peterson and Co., Evanstone, Ill., Elmford, N.Y. 1961
6. Coxeter H.S.M., Introduction to geometry, John Willey and Sons, Inc., N.Y.-London, 1961.
7. Dirac P.A.M., General theory of relativity, John Wiley and Sons, N.Y.- Sydney-Toronto, 1975
8. Bell I.S., Rev.Mod.Phys. 1966. V.38. P.447
9. von J.V.Neumann., Mathematische grundlagen der quantenmechanik, Verlag von Julius Springer, Berlin, 1932.